ORGANIC / SOFT CHEMICAL TREATMENTS

DOSAGE

Two insert strips per ten frames

PROPER PLACEMENT OF MEDICATION

Strips should be hung between brood frames

TREATMENT DURATION

• For best results, leave strip in the hive for a minimum of 14 days

HOW IT KILLS MITES

- Manufacturer's reported efficacy rate: 92% 94% (3 consecutive applications)
- Contact kills phoretic mites
- Bees walk on strips, chew strips, & disperse medication

RECOMMENDED APPLICATION TIME

- Temp Range for Application: Not temperature dependent
- Most effective when used during the pre-pollination period (before sealed brood), mid-summer and at the onset of winter brood development
- Use 3 consecutive applications during high brood development

USAGE WITH HONEY SUPERS

• Can be used with honey supers, however, we do not recommend

Manufacturer:

Beta Tec

Hopguard 3™

Active Ingredient:

www.betatec.com

Hop beta acids resin (16%)

PROS

- Not temperature dependent
- No adverse side effects to bees, brood, or queen
- · No residue in honey or wax

CONS

• Usually 3 consecutive treatments are needed to be effective



NOT FOR SALE. FOR EDUCATIONAL PURPOSES ONLY. © 2002-2023 EVERSWEET APIARIES. ALL RIGHTS RESERVED.

🦎 EVERSWEETAPIARIES.COM 🞉